
























POISONOUS PLANTS IN IOWA THAT AFFECT HORSES

COURTESY: PEGGY MILLER-AUWERDA, PHD - IOWA STATE UNIVERSITY DEPARTMENT OF ANIMAL SCIENCE

PLANT	TOXICITY	CLINICAL SYMPTOMS	PLANT	TOXICITY	CLINICAL SYMPTOMS
Alsike clover 	Low		Bristle grasses, foxtails 		Irritation and trauma to the mouth if ingested. Eye injuries.
Arrow, pod or goose grass 	Low	Tremors, frequent urination/defecation, gasping, convulsions	Buttercup 	Variable	Diarrhea, salivation
Azalea/laurel/rhododendron 		Acute colic, diarrhea, excessive salivation, depression, incoordination, stupor	Elderberry 	High	Tremors, frequent urination/defecation, gasping, convulsions
Black locust		Diarrhea or constipation, appetite loss, stupor, laminitis	Horsetail, maretail, horserush, or snake grass 	Low	Reluctance to move, diarrhea, constipation, possible blindness
Black walnut 	High	Edema in legs, increased heart/respiratory rate, increased temperature, laminitis	Japanese Yew 	Very high	Nervousness, difficult breathing, incoordination, convulsions
Bracken fern 	Low	Emaciation, incoordination, depression, paralysis, diarrhea, salivation. Found in forested areas.	Jimsonweed 		Excitability then depression, dilation of pupils, colic, diarrhea

<p>Locoweeds</p> 	<p>High</p>	<p>Incoordination, depression, shallow breathing, unsteadiness, coma</p>	<p>Ragwort/groundsel/hound's-tongue</p> 		<p>Weakness, liver failure, incoordination, yellow mucous membranes</p>
<p>Milkweed</p> 	<p>High</p>	<p>Incoordination, depression, shallow breathing, unsteadiness, coma</p>	<p>Red maple</p> 	<p>High</p>	<p>Dark/discolored mucous membranes, brown/red urine, depression, colic</p>
<p>Nightshade (Nightshade, potato and tomato)</p> 		<p>Trembling, incoordination, diarrhea</p>	<p>Saint-John's-wort</p> 		<p>Photosensitization</p>
<p>Oak (Tannins in leaves, bark or acorns, especially when green)</p>	<p>Variable</p>	<p>Colic, constipation followed by blood diarrhea, frequent urination, anorexia, depression</p>	<p>Sudan grass</p> 	<p>Low</p>	<p>Tremors, frequent urination/defecation, gasping, convulsions</p>
<p>Oats, Cultivated</p>		<p>Breathing difficulty, skin irritation, paralysis, convulsions, death (rarely).</p>	<p>Tall Fescue</p> 		<p>Reproductive problems, "poor doers", lameness, dry gangrene, fever, and death.</p>
<p>Onions, wild & domestic</p> 		<p>Weak, increased heart rate, increased respiration rate, mild to severe anemia, dark coffee colored urine, death</p>	<p>Thistles</p> 		<p>Irritation and trauma to the mouth if ingested. Eye injuries.</p>
<p>Poison hemlock</p> 	<p>High</p>	<p>Trembling, incoordination, salivation, colic, shallow breathing, coma, diarrhea</p>	<p>Water hemlock</p> 	<p>Very high</p>	<p>Violent spasms, rapid respiration and heart rate, coma</p>

REFERENCES:

Cornell University Poisonous Plants Web Page [<http://www.ansci.cornell.edu/plants/plants.html>]
 "Indiana Plants Poisonous to Livestock and Pets" [<http://vet.purdue.edu/depts/addl/toxic/cover1.htm>]
 "University of Illinois Plants Toxic to Animals" [<http://www.library.uiuc.edu/vex/toxic/intro.htm>]